
7-3 Cell Boundaries





Slide 1 of 47

7-3 Cell Boundaries Cell Membrane



What is the main function of the cell membrane?



Cell Membrane




The cell membrane regulates what enters and leaves the cell and also provides protection and support.



7-3 Cell Boundaries Cell Membrane

Cell Membrane






Slide 4 of 47

7-3 Cell Boundaries — Cell Membrane

The composition of nearly all cell membranes is a double-layered sheet called a **lipid bilayer**.






Slide 5 of 47

7-3 Cell Boundaries — Cell Membrane

Most cell membranes contain protein molecules embedded in the lipid bilayer, some of which have carbohydrate molecules attached to them.





Slide 6 of 47

7-3 Cell Boundaries Cell Walls



What is the main function of the cell wall?



7-3 Cell Boundaries Cell Walls



The main function of the cell wall is to provide support and protection for the cell.



Cell Walls

Cell walls are found in plants, algae, fungi, and many prokaryotes.

The cell wall lies outside the cell membrane.

Most cell walls are porous enough to allow water, oxygen, carbon dioxide, and certain other substances to pass through easily.



Diffusion Through Cell Boundaries

Every living cell exists in a liquid environment.

The cell membrane regulates movement of dissolved molecules from the liquid on one side of the membrane to the liquid on the other side.



Measuring Concentration

A solution is a mixture of two or more substances.

The substances dissolved in the solution are called solutes.

The **concentration** of a solution is the mass of solute in a given volume of solution, or mass/volume.





What happens during diffusion?



Diffusion


Particles in a solution tend to move from an area where they are more concentrated to an area where they are less concentrated.

This process is called **diffusion**.


When the concentration of the solute is the same throughout a system, the system has reached equilibrium.












There is a higher concentration of solute on one side of the membrane as compared to the other side of the membrane.








Solute particles move from the side of the membrane with a higher concentration of solute to the side of the membrane with a lower concentration of solute. The solute particles will continue to diffuse across the membrane until equilibrium is reached.








When equilibrium is reached, solute particles continue to diffuse across the membrane in both directions.







Diffusion depends upon random particle movements. Therefore, substances diffuse across membranes without requiring the cell to use energy.









Osmosis




Osmosis is the diffusion of water through a selectively permeable membrane.





How Osmosis Works





Slide 21 of 47

Water tends to diffuse from a highly concentrated region to a less concentrated region.

If you compare two solutions, the more concentrated solution is **hypertonic** ("above strength").

The more dilute solution is **hypotonic** ("below strength").



When concentrations of solutions are the same on both sides of a membrane, the solutions are **isotonic** ("same strength").



Osmotic Pressure

Osmosis exerts a pressure known as osmotic pressure on the hypertonic side of a selectively permeable membrane.




Because the cell is filled with salts, sugars, proteins, and other molecules, it will almost always be hypertonic to fresh water.

If so, the osmotic pressure should produce a net movement of water into the cell. As a result, the volume of the cell will increase until the cell becomes swollen or bursts.



Cells in large organisms are not in danger of bursting because they are bathed in fluids, such as blood, that are isotonic.

Other cells are surrounded by tough cell walls that prevent the cells from expanding even under tremendous osmotic pressure.





Facilitated Diffusion

Cell membranes have protein channels that act as carriers, making it easy for certain molecules to cross.



7-3 Cell Boundaries Facilitated Diffusion


The movement of specific molecules across cell membranes through protein channels is known as **facilitated diffusion**.

Hundreds of different protein channels have been found that allow particular substances to cross different membranes.



7-3 Cell Boundaries Facilitated Diffusion





Slide 29 of 47


Glucose



7-3 Cell Boundaries Facilitated Diffusion

Although facilitated diffusion is fast and specific, it is still diffusion.

Therefore, facilitated diffusion will only occur if there is a higher concentration of the particular molecules on one side of a cell membrane as compared to the other side.





Active Transport

Sometimes cells move materials in the opposite direction from which the materials would normally move—that is against a concentration difference. This process is known as **active transport**.

Active transport requires energy.



Molecular Transport


In active transport, small molecules and ions are carried across membranes by proteins in the membrane.

Energy use in these systems enables cells to concentrate substances in a particular location, even when diffusion might move them in the opposite direction.



active art

Active Transport





Endocytosis and Exocytosis

Large molecules and even solid clumps of material may undergo active transport by means of the cell membrane.


Endocytosis is the process of taking material into the cell by means of infoldings, or pockets, of the cell membrane.

The pocket breaks loose from the outer portion of the cell membrane and forms a vacuole within the cytoplasm.



Two examples of endocytosis are:

- phagocytosis
- pinocytosis





In **phagocytosis**, extensions of cytoplasm surround a particle and package it within a food vacuole. The cell then engulfs it.

Phagocytosis requires a considerable amount of energy.



In **pinocytosis**, tiny pockets form along the cell membrane, fill with liquid, and pinch off to form vacuoles within the cell.



Exocytosis

Many cells also release large amounts of material from the cell, in a process called exocytosis.

During **exocytosis**, the membrane of the vacuole surrounding the material fuses with the cell membrane, forcing the contents out of the cell.



Continue to:

Section QUIZ

- or -

Click to Launch:





- Unlike a cell wall, a cell membrane
- A a. is composed of a lipid bilayer.
 - b. provides rigid support for the surrounding cell.
 - c. allows most small molecules and ions to pass through easily.
 - d. is found only in plants, fungi, algae, and many prokaryotes.



- The concentration of a solution is defined as thea. volume of solute in a given mass of solution.
- b. mass of solute in a given volume of solution.
 - c. mass of solution in a given volume of solute.
 - d. volume of solution in a given mass of solute.



- If a substance is more highly concentrated outside the cell than inside the cell and the substance can move through the cell membrane, the substance will
 - a. move by diffusion from inside the cell to outside.
 - b. remain in high concentration outside the cell.



- c. move by diffusion from outside to inside the cell.
- d. cause water to enter the cell by osmosis.



Slide 42 of 47

- The movement of materials in a cell against a concentration difference is called
 - a. facilitated diffusion.
- b. active transport.
 - c. osmosis.
 - d. diffusion.



- The process by which molecules diffuse across a membrane through protein channels is called
 - a. active transport.
 - b. endocytosis.
- C. facilitated diffusion.
 - d. osmosis.



END OF SECTION